

Lattices compatible with regular polytopes

Peter McMullen

University College London, Gower Street, London WC1E 6BT, England, United Kingdom

Available online 11 March 2008

For the 80th birthday of Ludwig Danzer

Abstract

In a recent paper, Karpenkov has classified the lattice polytopes (that is, with vertices in the integer lattice \mathbb{Z}^d) which are regular with respect to those affinities which preserve the lattice. An alternative approach is adopted in this paper. For each regular polytope P in euclidean space \mathbb{E}^d , those lattices Λ are classified which are compatible with P , in the sense that some translate of Λ contains the vertices of P , and this translate is preserved by the symmetries of P .

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

A convex d -polytope P whose vertex-set $\text{vert } P$ is a subset of the integer lattice \mathbb{Z}^d in euclidean space \mathbb{E}^d is called *lattice regular* if it is regular with respect to the group \mathcal{U}_d of affinities which preserve \mathbb{Z}^d . Recall that being *regular* means that the subgroup $\mathcal{L}(P)$ of \mathcal{U}_d which preserves P is transitive on the family $\mathcal{F}(P)$ of *flags* of P , which are chains $F_0 < F_1 < \dots < F_{d-1}$ of faces of P . Karpenkov [2] has recently classified such lattice regular polytopes, up to equivalence under \mathcal{U}_d .

Here, we pursue an alternative approach to this problem. We start with a convex polytope P which is regular in the usual sense, with respect to the group \mathcal{I}_d of isometries of \mathbb{E}^d , and ask which lattices Λ are *compatible* with P , meaning that (up to translation) $\text{vert } P \subseteq \Lambda$, and the subgroup $\mathcal{G}(P)$ of \mathcal{I}_d of (isometric) symmetries of P preserves Λ . The advantage of looking at things from this viewpoint is that we no longer have to concern ourselves with the non-uniqueness arising from the equivalence under \mathcal{U}_d in the original problem.

E-mail address: p.mcmullen@ucl.ac.uk.

The outline of the rest of the paper is as follows. In Section 2, we consider the problem in general terms, in particular introducing the edge-module $\Lambda_e = \Lambda_e(P)$ of a polytope P . In Section 3, we eliminate the non-crystallographic cases, where Λ_e is non-discrete. In the following Sections 4–6, we treat the three infinite families of regular polytopes, namely, the simplices, cubes and cross-polytopes. In Section 7, we conclude the classification by dealing with the regular hexagon and 24-cell. Finally, we summarize the results in Section 8.

General references for the background to regular polytope theory are [1,6]; we shall usually follow the notational conventions of the latter.

2. General considerations

We begin by remarking that our rephrasing of the problem is equivalent to that of Karpenkov [2]. A finite group of affinities (such as a subgroup of \mathcal{U}_d) is conjugate under a suitable affinity Φ to a group of isometries, and so, if a convex polytope P is lattice regular, then some affine image ΦP of P is actually regular in the usual sense (compare [3]), and the corresponding image $\Lambda = \Phi \mathbb{Z}^d$ of \mathbb{Z}^d is a lattice which is compatible with ΦP .

Conversely, if we are given a regular polytope P in \mathbb{E}^d and a lattice Λ compatible with P , then we can apply an affinity Ψ taking a basis of Λ into the standard basis of \mathbb{E}^d (and thus Λ into $\Psi\Lambda = \mathbb{Z}^d$), so that ΨP will be lattice regular in the sense of [2].

We thus start with a convex d -polytope P in \mathbb{E}^d which is regular in the usual sense, under the group $\mathcal{G} = \mathcal{G}(P)$ of isometries which preserve P . (We shall always make the assumption that polytopes have dimension d , and lattices have rank d .) As in Section 1, we say that a lattice Λ in \mathbb{E}^d is *compatible* with P if there is some translate $\Lambda' = \Lambda + t$ of Λ such that $\text{vert} P \subseteq \Lambda'$ and $\mathcal{G}\Lambda' = \Lambda'$. (We phrase things this way, because it will sometimes be convenient not to have the vertices of P sitting in an actual lattice.)

The *edge-module* $\Lambda_e = \Lambda_e(P)$ is defined as follows: if

$$\mathcal{E} = \mathcal{E}(P) := \{v - w \mid v, w \in \text{vert } P\},$$

then $\Lambda_e := \mathbb{Z}[\mathcal{E}]$ is the \mathbb{Z} -module generated by \mathcal{E} . In the definition of Λ_e , it is clear that we need only take as generators the *edge-vectors* $v - w$, where v, w are the two vertices of an edge of P ; this accounts for the nomenclature. If Λ, Λ' are lattices such that Λ is a sublattice of Λ' , then we say that Λ' is a *superlattice* of Λ . We first have an obvious result.

Theorem 2.1. *If Λ is a lattice which is compatible with the regular polytope P , then Λ is a superlattice of $\Lambda_e(P)$.*

Proof. Indeed, Λ must contain all edge-vectors in $\mathcal{E}(P)$, and the claim follows at once. \square

Remark 2.2. Of course, there are extra conditions imposed on such a superlattice Λ by the symmetries of P .

Recall that a vector v in a lattice Λ is *primitive* if $v \neq kw$ for any $w \in \Lambda$ and (integer) $k \geq 2$. We call a lattice Λ which is compatible with a regular polytope P *primitive* if each edge-vector in $\mathcal{E}(P)$ is primitive in Λ . If $q \in \mathbb{N}$, then

$$\frac{1}{q}\Lambda := \left\{ \frac{1}{q}v \mid v \in \Lambda \right\}$$

has the obvious meaning. It is clear that a lattice Λ which is compatible with P is of the form $\Lambda = \frac{1}{q}\Lambda'$ for some $q \in \mathbb{N}$ and some primitive compatible lattice Λ' ; thus it suffices to determine the primitive lattices.

The subsequent discussion is aided by the following remark.

Proposition 2.3. *Let Λ be a lattice in \mathbb{E}^d for some d , and let Λ' be a superlattice of Λ . Then Λ' is a sublattice of $\frac{1}{q}\Lambda$ for some q .*

Proof. A basis of Λ consists of integer linear combinations of vectors of a basis of Λ' . It follows at once that a basis of Λ' consists of rational linear combinations of vectors of a basis of Λ . The conclusion of the proposition is immediate. \square

Last in this section, we recall that the *reciprocal lattice* of Λ is

$$\Lambda^* = \{y \in \mathbb{E}^d \mid \langle x, y \rangle \in \mathbb{Z} \text{ for all } x \in \Lambda\}.$$

Note that $\Lambda^* \cong \text{Hom}(\Lambda, \mathbb{Z})$ (as abelian groups). We shall see that, if Λ is compatible with some regular polytope P , then a suitable multiple of Λ^* will be as well.

3. The non-crystallographic cases

The edge-module Λ_e of a regular polytope P is generated by finitely many vectors, and so is of finite rank over \mathbb{Z} . However, it is not necessarily a lattice, by which we mean that it is discrete. In a more general context (see [4]), we have called polytopes for which Λ_e is discrete *rational*; in the convex case, the polytopes are usually referred to as *crystallographic*. In this section, we dispose of the non-crystallographic cases.

We begin with the regular polygons $\{p\}$ for $p \geq 3$.

Theorem 3.1. *The edge-module $\Lambda_e(P)$ of a regular p -gon P is a lattice if and only if $p = 3, 4$ or 6 .*

Proof. If Λ_e is a lattice in \mathbb{E}^2 , then it has a shortest vector v , say. We lose no generality in scaling so that $\|v\| = 1$. Let Φ_p denote rotation about the origin o through the angle $2\pi/p$, so that Φ_p is a symmetry of Λ_e . For $p \geq 7$, we have

$$\|\Phi_p v - v\| = 2 \sin \frac{\pi}{p} < 1,$$

contradicting the definition of v . For $p = 5$, we have

$$\|\Phi_5^2 v + v\| = 2 \sin \frac{\pi}{10} < 1,$$

again a contradiction. Thus all cases except for $p = 3, 4$ or 6 have been excluded; that these are crystallographic is well known, and will in any event be dealt with later. \square

The remaining non-crystallographic regular polytopes are those with 5-fold symmetries.

Theorem 3.2. *The edge-module $\Lambda_e(P)$ is non-discrete if P is a dodecahedron, icosahedron, 120-cell or 600-cell.*

Proof. Each of the dodecahedron and 120-cell has (2-dimensional) faces which are regular pentagons. The links of a vertex of an icosahedron and edge of a 600-cell are also regular

pentagons. Thus, in every case, the edge-module contains a submodule congruent to the edge-module of a regular pentagon; this is non-discrete by [Theorem 3.1](#), which implies the claim of the theorem. \square

4. Simplices

The case of the regular d -simplex T is most easily treated in \mathbb{E}^{d+1} . We choose the vertices of T to be the standard basis vectors e_0, \dots, e_d of \mathbb{E}^{d+1} . The symmetry group $\mathcal{G} = \mathcal{G}(T)$ of T thus consists of all permutations of e_0, \dots, e_d , and so of all permutations of the coordinates of \mathbb{E}^{d+1} . We see at once that we have

Theorem 4.1. *The edge-module $\Lambda_e(T)$ of the regular d -simplex T is*

$$\begin{aligned}\Lambda_1 &:= \langle e_i - e_j \mid 0 \leq i < j \leq d \rangle \\ &= \{(\xi_0, \dots, \xi_d) \in \mathbb{Z}^{d+1} \mid \xi_0 + \dots + \xi_d = 0\}.\end{aligned}$$

Ignoring the scaling factor $1/q$ in the previous discussion, our next task is to determine the sublattices Λ of $\Lambda_e = \Lambda_1$ which have the symmetry of T . Our arguments exactly parallel those of [\[5\]](#); see also [\[6, Sections 6D, 6E\]](#). Let s be the smallest positive difference $\alpha_i - \alpha_j$ which occurs in vectors $a = (\alpha_0, \dots, \alpha_d) \in \Lambda$. Permuting coordinates as necessary, we can assume that $i = 0$, $j = 1$ and that $s = \alpha_0 - \alpha_1$. If a' is obtained from a by the transposition $(0\ 1)$ which interchanges the first two coordinates, then

$$a - a' = (s, -s, 0^{d-1}),$$

where r^k in a vector denotes a string r, \dots, r of length k . Again applying the symmetries of T , we see that $s\Lambda_1$ is a sublattice of Λ .

We now add to a suitable multiples of the vectors $s(e_0 - e_j)$, so as to ensure that the (new) coordinates α_j satisfy $-s < \alpha_j \leq 0$ for $j = 1, \dots, d$. It is immediate that all such α_j are equal; otherwise, we have a smaller difference than s between two coordinates of a vector in Λ . If all these $\alpha_j = 0$ (with all possible choices of a), then $\Lambda = s\Lambda_1$; bear in mind that Λ is a sublattice of Λ_1 . Alternatively, Λ contains a vector rb for some $r \in \mathbb{N}$, where $b = (d, (-1)^d)$ (with the convention introduced before). In fact, if we write Λ_b for the sublattice of Λ_1 generated by b and its images under the symmetries of T , then we actually have $\Lambda \geq r\Lambda_b$, and $s = (d+1)r$. We further note that, if $b' := (-1, d, (-1)^{d-1})$, then $b' \in \Lambda_b$, and $b - b' = (d+1)(1, -1, 0^{d-1})$; we conclude that

$$(d+1)\Lambda_1 \leq \Lambda_b. \tag{4.1}$$

Now let us revert to the original problem. A lattice Λ which is compatible with T contains $\frac{1}{q}\Lambda_1$ for some $q \in \mathbb{N}$ (q is not fractional, because Λ contains Λ_1), and possibly some $r\Lambda_b$ with a suitable fractional r . If Λ is primitive, then $q = 1$, and we can write

$$\Lambda = r\Lambda_b \quad \text{or} \quad \Lambda_1 + r\Lambda_b$$

for some (positive) rational r (recall [\(4.1\)](#)). If $r = p/q$ in its lowest terms (not the same q as before), then $kp \equiv 1 \pmod{q}$ for some $k \in \mathbb{N}$, and since $kr\Lambda_b \leq r\Lambda_b$ and $\Lambda_b \leq \Lambda_1$, we can therefore assume that $r = 1/q$ for some $q \in \mathbb{N}$. Moreover, by [\(4.1\)](#), it follows that $(d+1)r\Lambda_1 = r\Lambda_b \leq \Lambda$. Hence, if Λ is to be primitive, then $(d+1)r$ cannot be fractional; it follows that $(d+1)r = (d+1)/q \in \mathbb{N}$, so that q is a divisor of $d+1$.

Summarizing this discussion, we see that we have proved

Theorem 4.2. *The primitive lattices which are compatible with the regular d -simplex T are*

$$\begin{cases} \Lambda_1, \\ \Lambda_q := \Lambda_1 + \frac{1}{q} \Lambda_b, & \text{for some } q|(d+1) \text{ with } 1 < q < d+1, \\ \Lambda_{d+1} := \frac{1}{d+1} \Lambda_b. \end{cases}$$

There is a comparatively easy way of getting at the indices of $\Lambda_e = \Lambda_1$ in these superlattices. Define

$$\Gamma_k := \{(\xi_0, \dots, \xi_d) \in \mathbb{Z}^{d+1} \mid \xi_0 + \dots + \xi_d = k\};$$

thus $\Lambda_1 = \Gamma_0$. Then it is clear that

$$\Gamma_k + \Gamma_m = \Gamma_{k+m}$$

for each k and m . Next, define $H_k := \text{aff} \Gamma_k$ to be the hyperplane containing Γ_k . If Π denotes the orthogonal projection onto H_0 , then $\Pi \Gamma_{d+1} = \Lambda_1$. Further, we see that

$$\Pi e_0 = \frac{1}{d+1} b,$$

whose images under \mathcal{G} generate Λ_{d+1} . It is then an easy matter to conclude that

$$[\Lambda_q : \Lambda_1] = q$$

for each divisor q of $d+1$.

We end the section by observing that $\Lambda_{d+1} = \Lambda_1^*$. Indeed, we actually have

Proposition 4.3. *The primitive lattices compatible with the regular simplex occur in reciprocal pairs:*

$$\Lambda_q^* = \Lambda_{(d+1)/q}$$

for each $q|(d+1)$.

5. Cubes

For the d -cube C , the procedure is more straightforward than that of Section 4. The standard unit cube has vertex-set $\{0, 1\}^d$, and we see at once that

$$\Lambda_e = \Lambda_1 := \mathbb{Z}^d.$$

The group $\mathcal{G} = \mathcal{G}(C)$ of C consists (up to translations) of all permutations of the coordinates of \mathbb{E}^d , together with all changes of their signs. We employ similar arguments to those of Section 4 (see [5] or [6, Section 6D]), but for completeness it is worth briefly recalling the details here.

We are thus looking for sublattices of \mathbb{Z}^d on which \mathcal{G} acts; we may assume that $d \geq 2$. Let $\lambda \in \mathbb{Z}^d$; in the present case, we let $a = (\alpha_1, \dots, \alpha_d) \in \Lambda$ be a vector with smallest positive coordinate $s = \alpha_j$. Under the action of \mathcal{G} , we can suppose that $j = 1$; changing the sign of the first coordinate and subtracting the new vector from the old shows that $(2s, 0^{d-1}) \in \Lambda$. Reducing each coordinate of a modulo $2s$ now shows that Λ is generated by the vector $(s^k, 0^{d-k})$ and its

images under \mathcal{G} , for some $k \geq 1$. However, if $3 \leq k < d$, then $(0, s^k, 0^{d-k-1}) \in \Lambda$ also, which leads quickly to $(s, s, 0^{d-2}) \in \Lambda$ (and then, if k is odd, to $(s, 0^{d-1}) \in \Lambda$ as well).

In conclusion, we have shown that the sublattices of Λ_1 which are preserved by \mathcal{G} are $s\Lambda_1$, $s\Lambda_2$ or $s\Lambda_d$, generated respectively by

$$s(1, 0^{d-1}), \quad s(1, 1, 0^{d-2}) \quad \text{or} \quad s(1^d) \quad (5.1)$$

and their images under \mathcal{G} , for some $s \in \mathbb{N}$ (of course, $\Lambda_2 = \Lambda_d$ when $d = 2$; for convenience, we have changed the notation from [5,6]). Apart from the obvious subgroup relationships, we have

$$\begin{aligned} \Lambda_2 &\leq \Lambda_1, & \text{with index } [\Lambda_1 : \Lambda_2] &= 2, \\ \Lambda_d &\leq \Lambda_1, & \text{with index } [\Lambda_1 : \Lambda_d] &= 2^{d-1}, \end{aligned}$$

and, if d is even,

$$\Lambda_d \leq \Lambda_2, \quad \text{with index } [\Lambda_2 : \Lambda_d] = 2^{d-2}.$$

Further, we note that

$$[\Lambda_2 : 2\Lambda_1] = 2^{d-1}, \quad [\Lambda_d : 2\Lambda_1] = 2.$$

Finally, Λ_1 is self-reciprocal, while

$$\Lambda_2^* = \frac{1}{2}\Lambda_d.$$

It is now an easy matter to deduce

Theorem 5.1. *The primitive lattices which are compatible with the regular d -cube C are*

$$\left\{ \begin{array}{l} \Lambda_1, \\ \frac{1}{2}\Lambda_d, \\ \frac{1}{2}\Lambda_2. \end{array} \right.$$

6. Cross-polytopes

The analysis for the regular d -cross-polytope X is very similar to that for the cube. Taking the vertices of X to be all $\pm e_j$ for $j = 1, \dots, d$, with $\{e_1, \dots, e_d\}$ the standard basis of \mathbb{E}^d , we see at once that $\Lambda_e = \Lambda_e(X) = \Lambda_2$ in the notation of Section 5. We have already detailed the subgroup relationships in that section (it is helpful to bear in mind that $\Lambda_2 \leq \Lambda_1$, and so the previous analysis applies), and the main result follows quickly.

Theorem 6.1. *The primitive lattices which are compatible with the regular d -cross-polytope X are*

$$\left\{ \begin{array}{l} \Lambda_2, \\ \Lambda_1, \\ \frac{1}{2}\Lambda_d, \end{array} \right.$$

with Λ_r as defined by (5.1).

7. The anomalous cases

There remain two polytopes to be considered: these are the hexagon and the 24-cell.

We can treat the hexagon H most easily by observing that it is the Minkowski (vector) sum $T - T$ of a regular triangle T and its opposite $-T$. This tells us at once that its edge-module $\Lambda_e(H)$ is the same as that of T , namely, Λ_1 in the notation of Section 4 (with $d = 2$). The only other divisor of $2 + 1 = 3$ is 3, and we quickly conclude that

Theorem 7.1. *The primitive lattices which are compatible with the regular hexagon H are the reciprocal lattices*

$$\left\{ \begin{array}{l} \Lambda_1, \\ \frac{1}{3}\Lambda_b, \end{array} \right.$$

in the notation of Section 4.

Remark 7.2. Note that the centre of the hexagon H is a point of $\Lambda_e(H)$.

We finally come to the 24-cell V . For simplicity of notation, we take its vertices to be

$$\pm 2e_j \quad \text{for } j = 1, \dots, 4, \quad (\pm 1, \pm 1, \pm 1, \pm 1),$$

with $\{e_1, \dots, e_4\}$ the standard basis of \mathbb{E}^4 . Then its edge-module is Λ_4 in the notation of Section 5. Now Λ_1 does not have the symmetry of V , whereas Λ_2 does; the latter is the edge-module of the 24-cell with vertices

$$\pm e_j \pm e_k \quad \text{for } 1 \leq j < k \leq 4.$$

Note that Λ_4 is similar to Λ_2 , but $\sqrt{2}$ times as large. Comparing [5] or [6, Section 6E] shows that (up to scaling) these two lattices are the only ones with the symmetry of V , and thence it is straightforward to deduce

Theorem 7.3. *The primitive lattices which are compatible with the regular 24-cell V are*

$$\left\{ \begin{array}{l} \Lambda_4, \\ \Lambda_2, \end{array} \right.$$

with Λ_r as defined by (5.1) with $d = 4$.

Remark 7.4. Note that the 24-cell V has diametral hexagons, and hence the centre of V is a point of $\Lambda_e(V)$. We have already observed that Λ_4 and $\frac{1}{2}\Lambda_2$ are reciprocal lattices.

8. Summary of results

We end with a table which summarizes the results obtained in the paper. Here, $\ell(P)$ is the number of primitive lattices which are compatible with the regular polytope P , and $\delta(k)$ denotes the number of divisors of $k \in \mathbb{N}$.

d	polytope P	$\ell(P)$
1	segment	1
2	triangle	2
	square	2
	hexagon	2
$d \geq 3$	simplex	$\delta(d + 1)$
	cube	3
	cross-polytope	3
4	24-cell	2

Acknowledgements

We are grateful to Oleg Karpenkov for an early sight of his paper [2], and for some helpful correspondence. Following a useful suggestion of Barry Monson, we expanded the discussion of the sublattices of \mathbb{Z}^d in Section 5. Egon Schulte pointed out several errors in an earlier draft, and made further valuable comments on it. Finally, thanks are due to the referees for catching several further mistakes.

References

[1] H.S.M. Coxeter, Regular Polytopes, 3rd edition, Dover, New York, 1973.
[2] O. Karpenkov, Classification of lattice-regular lattice convex polytopes, *Funct. Anal. Other Math.* 1 (2006) 19–42.
[3] P. McMullen, Affine and projectively regular polytopes, *J. London Math. Soc.* 43 (1968) 755–757.
[4] P. McMullen, Regular polytopes of full rank, *Discrete Comput. Geom.* 32 (2004) 1–35.
[5] P. McMullen, E. Schulte, Higher toroidal regular polytopes, *Adv. Math.* 117 (1996) 17–51.
[6] P. McMullen, E. Schulte, Abstract Regular Polytopes, in: *Encyclopedia of Mathematics and its Applications*, vol. 92, Cambridge University Press, 2002.